

The origin of cores and density profiles of gaseous baryonic structures

Steen H. Hansen & Joachim Stadel

University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland

ABSTRACT

We study the origin of cores and density profiles of gaseous baryonic structures in cosmology. By treating the baryons as a viscous gas, we find that both spheres and disks are possible solutions. We find analytically that the density profiles have inner and outer solutions, which in general are different. For disks we identify a central core, with density profile $\rho_d = \text{constant}$, and the outer profile $\rho_d \sim r^{-3}$. For spherical structures we find the profile $\rho_s \sim r^{-6}$. In the presence of a dominating central black hole we find the inner profile $\rho \sim r^{-3/2}$. When the mass is dominated by a dark matter component then the baryonic density profile will depend on the dark matter profile, and we point out how one can use this connection to infer the DM profile directly by observing the baryonic density profile.

Subject headings: galaxies: clusters: general — galaxies: structure — methods: analytical

1. Introduction

The universe is full of large gaseous baryonic structures such as galaxies and clusters. These structures are observed to take on a variety of shapes ranging from disks to spherical configurations. The radial profiles of these structures have been observed to be rather non-trivial, e.g. changing from one power-law profile near the centre to another power-law profile at large radii. Even though these baryonic configurations have been observed and studied for many years, there is surprisingly little theoretical guidance to understand the origin of such complexity. Astronomers have for years been using phenomenological profiles, ones with differing inner and outer shape, however, no simple theoretical explanation for this structure exists.

In recent years numerical analyses have improved enormously, and we can now simulate much of the structures we observe. Never the less, it is important to have a simple understanding of the underlying physics, which can be obtained most easily through analytical studies of the basic equations.

In this letter we attempt an analytical treat-

ment of the basic equations, asking which density profiles are expected of purely gaseous baryonic structures, and which profiles lead to stable baryonic structures within dark matter halos. We address these questions by considering a fluid approach whereby we analyse asymptotic stable solutions of the Navier-Stokes equations. This approach is new in astrophysics and surprisingly simple. Our main results are, that the density profile of gaseous baryonic structures is fairly complex, namely that the radial density profile may have different slopes in the inner and outer regions. This is an issue which has been much discussed for Dark Matter profiles, but which has not been understood previously for baryonic structures. We find that spheres and disks are the only possible solutions. From an observational point of view it seems obvious that these are possible solutions, and while the creation of disks is understood theoretically, the fact that no other solutions are stable is non-trivial. As a byproduct of our analysis we show how one can infer the DM density profile purely by observing the baryonic density profile. We also analyse how a central massive black hole influences the solutions.

2. Solving Navier-Stokes equations

The behaviour of any collisional gas or fluid is fully determined by the Navier-Stokes (N-S) equations, which are 3 hydrodynamical equations for the velocity vector, and a continuity equation. In the following we will make several assumptions, which certainly limits the applicability of our results, however, the solutions we find should be valid for the description of general properties of structures such as gaseous baryonic galaxies and haloes. The main assumption is that there are sufficient collisions while the structure forms. Thus if a gaseous baryonic structure has formed and subsequently all the baryons condense into stars, then the star density profile should still approximately follow the original density profile. Baryons often have sufficient collisions to equilibrate, e.g. in a typical intra-cluster gas the equilibration timescale is about 10^7 yrs, with mean free path of tens of kpc compared to radii of few Mpc. We do not expect that our findings should apply to dark matter structures, since dark matter presumably does not have sufficient collisions to ensure the validity of the N-S equations.

In spherical coordinates the N-S equations describe the velocity vector (v_r, v_Θ, v_ϕ) . Here r is the radial coordinate, Θ is the angle in the xy -plane, and ϕ is the angle from the z -axis. The form of the equations is well known, see e.g. Landau & Lifshitz (1987), and contains time-derivatives, pressure terms, viscosity terms and a gravitational term.

Our first 2 assumption are, that the gas has reached a stable configuration (no time-derivative), and that it has picked out an orientation in space, in such a way that all the gas is moving only in the Θ -direction. Thus we have $v_r = 0$ (no contraction or expansion), and $v_\phi = 0$. Here one must keep in mind, that by considering the N-S equations we are taking a fluid approach, which implies that we are following a fluid element, and this basically corresponds to averaging over all the particles moving through the fluid element. For the Θ -velocity we consider the general form

$$v_\Theta = v_\alpha \left(\frac{r}{r_\alpha} \right)^\alpha (\sin\phi)^\chi, \quad (1)$$

where α and χ are constants to be determined, v_α and r_α are unknown constants, with the physical

interpretation that r_α is a characteristic radius, and v_α is the velocity of the fluid element at that radius.

2.1. The Θ -equation

The v_Θ -equation becomes very simple with the assumed form of the velocities

$$0 = \nu \left[\nabla^2 v_\Theta - \frac{v_\Theta}{r^2 \sin^2 \phi} \right], \quad (2)$$

where ∇^2 is the scalar Laplacian, and ν is the kinematic viscosity. For now all that matters is the existence of a non-zero viscosity, so the absolute magnitude (and even radial dependence) is not important for the results¹. When we use the form for v_Θ in eq. (1), then equation (2) has 4 solutions

$$\chi = +1 \quad , \quad \alpha = 1, -2, \quad (3)$$

$$\chi = -1 \quad , \quad \alpha = 0, -1. \quad (4)$$

The solution with $\chi = +1$ is exactly what one should expect for spherical symmetry, and we will refer to these solutions as the '*spherical solutions*'. The solutions with $\chi = -1$ indicate unstable solutions everywhere except in the flat cylinder where $\sin\phi = 1$, and we will refer to these as '*disk solutions*'. We thus see that both spherical and disk solutions exist, and that they have different rotational structure (different α). Now, looking at eq. (1), it is clear, that the solutions with negative α are divergent for $r \rightarrow 0$, and we will therefore refer to those solutions as '*outer solutions*', and similarly, the solutions with positive α are inconsistent for large radii, and we will refer to those as '*inner solutions*'.

There are several obvious extensions of the method described here. One could in particular allow for a general Θ -dependence in v_Θ , a non-zero radial velocity, or an exponential radial dependence of velocities and density profile².

¹Certainly baryons have a non-zero viscosity, however the viscosity could also arise from turbulence in which case it could be very large, $\nu_{\text{turb}} \sim l\Delta v$ (Landau & Lifshitz 1987), where the dimension l is the size of the turbulent eddies, and Δv is the velocity dispersion.

²Allowing a Θ dependence in v_Θ as $v_\Theta \sim (\sin n\Theta)^\sigma$, with n the number of spiral arms, gives $\sigma = 0, 1$ for all n . Allowing for a non-zero radial velocity, $v_r \sim r^\tau$, gives $\tau = 1, -2$.

Already from this one sees that a spherical solution, in eq. (3), must have different radial structure for the inner and outer solutions (because the only solutions are $\alpha = 1, -2$), and hence one may expect to find different density profiles in the central and outer regions. We point out that such phenomenon of simultaneous existence of two flow patterns is rather common in hydrodynamics, the simplest may be the hydraulic jump which is observed as a several centimetre large circular ring in any kitchen sink, when the flowing water goes from a $1/r$ profile to a constant, see Hansen et al. (1997); Watanabe et al. (2003). We emphasize that a non-zero viscosity appears to be a necessary condition for the existence of these specific solutions, even though a fundamental understanding of how the microscopic physics (viscosity) can determine the macroscopic properties (the general flow pattern) is still generally missing in hydrodynamics.

2.2. The r -equation

From the Θ -equation, we identified the general flow pattern, and we noticed the possibility that one may have different velocity-flows in the inner and outer region. We will now use the v_r -equation to try to extract the asymptotic radial density profiles. Also the r -equation is very simple

$$-\frac{v_\Theta^2}{r} = -\frac{1}{\rho} \frac{\partial P}{\partial r} - \frac{M(r)G}{r^2}, \quad (5)$$

where ρ is the radially dependent density, P is the pressure, G is the gravitational constant, and $M(r)$ is the mass within the radius r . We assume that the pressure and density are related through $P = P_\alpha (\rho/\rho_\alpha)^\gamma$, where P_α and ρ_α are the unknown pressure and density at r_α . We assume the gas is monatomic with $\gamma = 5/3$. Let us consider densities of the form

$$\rho(r) = \rho_\alpha \left(\frac{r}{r_\alpha} \right)^\beta, \quad (6)$$

such that the parameter β determines the density profile. It is worth emphasising that it is exactly this β which we are trying to find.

Let us study the radial dependence of the 3 terms in eq. (5). Using v_Θ in eq. (1) the first (kinetic) term of eq. (5) goes like $v_\Theta^2/r \sim r^{2\alpha-1}$. The pressure gradient term goes like $1/\rho \cdot \partial P/\partial r \sim$

$r^{\delta\beta-1}$, where we have used $\delta = \gamma - 1 = 2/3$. The last (gravitational) term including $M(r)$, depends on the given system we are considering. If the mass is dominated by a point gravitational source (e.g. a central black hole), then it goes like $M(r)G/r^2 \sim r^{-2}$. If the mass is dominated by the matter density, then it goes like $M(r) \sim \int \rho(r)dV$, with dV the volume element. For spherical solutions this gravitational term thus goes like $r^{\beta+1}$ with β from eq. (6), and for disk solutions it goes like r^β . This gravitational term has the correct form for spherical distributions (and point sources), but is only an approximation for the pure disk case. Technically speaking the mass is logarithmic divergent for spherical structures with $\beta = -3$, however, the formula $M \sim r^{\beta+3}$ holds for any β arbitrarily close to -3 , and furthermore in a real situation there would be an outer cut-off. To be explicit, we are looking for solutions to an equation of the form

$$r^{2\alpha-1} = r^{\delta\beta-1} + r^\kappa, \quad (7)$$

where $\kappa = \beta + 1, \beta, -2$ for spherical, disk and BH matter dominance respectively. When we use the word 'solve' in the following, we are really just using the standard method of divergence cancellation, in the sense that the most divergent terms must cancel with each other. The optimal case is naturally that all divergences disappear, a case which we will refer to as 'good'. From our simple analysis the transition radius, r_α , which separates the inner from outer region, is not uniquely determined. We only find approximate disk relations like $r_\alpha \sim v_\alpha^2/G\rho_\alpha$. A full study including the coefficient is significantly more involved, and we will leave that for a later analysis³.

We want to solve eq. (5) for the profile parameter β , however, there are 4 situations to consider (the 4 different α from eqs. (3,4)), and for each case we can choose which is the dominating mass contribution, either from a sphere, from a disk or from a black hole (BH). The discussion in sections 2.3 and 2.4 is somewhat technical, and the reader is encouraged to look at figs. 1 and 2, which summarize the main findings of these sections.

³We note that when the v_Θ^2 term is negligible this eq. (5) is exactly the hydrostatic equilibrium, which is often written as $M(r) \sim rT\gamma\beta$. Furthermore, in the same notation one finds that the velocity dispersion, σ_v^2 , has a radial dependence, $d \ln \sigma_v^2 / d \ln r = \delta\beta$.

2.3. Disk

Let us first consider the $\alpha = 0$ case, which can be both an inner and outer solution. For the inner solution we find two density profiles, $\beta = 0, -3$. The $\beta = 0$ solution is more natural, because the first two divergent terms in eq. (5) (or similarly eq. (7)) can cancel, and the last term is non-divergent for $r \rightarrow 0$, whereas for the $\beta = -3$ solution, the two most divergent terms cancel, but the remaining term is still slightly divergent towards the centre. With this kind of argument one can divide all the solutions into ‘good’, and ‘reasonable’ solutions, and we will emphasise which are *good* solutions. If the mass is dominated by a BH, then the solution is $\beta = -3/2$. If the disk is dominated by a spherical distribution (either from a large fraction of the baryons, or from an unknown dark matter component) with profile β_s , then the solution is $\beta = 3/2(\beta_s + 2)$. In conclusion we see, that the only good solution for the inner disk is $\beta = 0$. In principle the $\alpha = 0$ case can also be an outer solution. In that case the solution is $\beta = -1$. If the mass is dominated by a BH, then $\beta = 0$ (good), and if dominated by a spherical distribution then $\beta \leq 0$.




Fig. 1.— A summary of disk solutions. The full lines (*self*, $\beta = 0, -3$) are where the mass is dominated by the disk itself. The dotted line (*BH*, $\beta = 0$) is for black hole domination, and the dashed line (*sphere*, $\beta = -3$) is for matter domination by a sphere with profile, $\beta_s = -3$.

The $\alpha = -1$ (which is an outer solution) gives $\beta = -3$ (good). If a BH dominates then $\beta = -3/2$, and if another spherical component domi-

nates (with profile β_s) then $\beta = 3/2(\beta_s + 2)$, where the case $\beta = -3$ is the only good solution. We summarize the good disk solutions in Figure 1.

To conclude the disk solutions, when the mass is dominated by the matter itself, then there is only 1 good configuration (we emphasise that by good we mean optimal removal of divergences), which has the inner slope of $\beta = 0$ and outer slope of $\beta = -3$. Following the similarity with the hydraulic jump in the kitchen sink, we will refer to such configuration as a galactic hydraulic drop.

2.4. Sphere

For the sphere it turns out that almost all the solutions are good, in the sense that all the divergences cancel in a simple manner. The $\alpha = 1$ is the inner solution, for which we find $\beta = 3, -6$. The $\beta = 3$ is obviously strange, and probably non-physical (a positive β would lead to a wrong sign in front of the pressure gradient term). With BH dominance one has $\beta = -3/2$, and if dominated by a disk (which could either be from a fraction of the same gas, or from another particle type) with profile β_d one finds $\beta = 3/2(\beta_d + 1)$. If dominated by another spherical distribution (which probably should arise from another particle type) with profile $\tilde{\beta}_s$, then we find $\beta = 3/2(\tilde{\beta}_s + 2)$.

The outer solution, $\alpha = -2$ gives $\beta = -6$. If a BH dominates then $\beta = -3/2$, and again if a disk (or sphere) dominates then $\beta = 3/2(\beta_d + 1)$ (or $3/2(\tilde{\beta}_s + 2)$). We summarize the good spherical solutions in Figure 2.

We note that if the mass is disk dominated with disk-profile $\beta_d = -3$, then the sphere happens to also get the profile $\beta_s = -3$. Thus one can envisage a significant fraction of the matter collapsing to a disk, which very well could take the profile $\beta_d = -3$ (as shown in section 2.3), and this would force the remaining matter, which is in a spherical configuration, to take the density profile $\beta_s = -3$. We also point out that if the mass is dominated by a dark matter sphere, then the baryon profile becomes $\beta = -3/2$ for a dark matter profile $\tilde{\beta}_s = -3$ as suggested by dark matter N-body simulations. If instead the dominating DM profile is $\tilde{\beta}_s > -2$, this would seemingly imply a positive baryonic slope, which is difficult to interpret. We feel that this limited applicability warns, that a more general calculation may provide a different




Fig. 2.— A summary of spherical solutions. The full lines (*self*, $\beta = -6$) are where the mass is dominated by the sphere itself, and the dotted lines (*BH*, $\beta = -3/2$) are for black hole domination. The dashed lines (*sphere*, $\beta = -3/2$) are for matter domination by another sphere with profile -3, and the dot-dashed lines (*disk*, $\beta = -3$) are for domination of a disk with profile -3.

connection between the DM and gaseous baryonic profiles

3. Discussion

An interesting possibility now appears, namely that one can use our results to infer the dark matter profiles from observations of the baryonic profile. This method can be used quite generally to infer the DM distribution, and is therefore complementary to other methods such as lensing observations. Let us say we have observed a baryonic sphere with profile β_{baryon} , and we know that the mass is dark matter dominated. Under the assumption that the DM is spherical we have $\beta_{\text{baryon}} = 3/2 (\tilde{\beta}_{\text{DM}} + 2)$. Thus, if we observe e.g. $\beta_{\text{baryon}} = -1.5$, then we know, that the DM has profile $\tilde{\beta}_{\text{DM}} = -3$. Determining the baryonic density profile directly, i.e. independent of dynamics, can be accomplished in several ways:

1) X-rays: observations of the luminosity, L_x , in various bands and in different radial bins gives the radial electron density of the plasma. Here the main concern is the validity of hydrostatic equilibrium and disentangling any cooling flow in the centre of the cluster. Relaxed clusters with no evidence of cooling flows do exist, for example A2029

(see Figure 1 of (Lewis et al. 2003)). The gas in this cluster shows an inner profile $\beta_{\text{baryon}} = -0.55$ and an outer $\beta_{\text{baryon}} = -1.62$ (note our differing definition of β here). The outer value is certainly DM dominated and our analysis implies that $\tilde{\beta}_{\text{DM}}$ is very close to -3 , which is expected from CDM simulations (Navarro et al. 1996; Moore et al. 1999). We do not attempt to deduce the inner DM slope here, because one cannot be certain that the mass is DM dominated at such small radii where the baryonic slope should reach its asymptotic value. Again we refer to the warning in the end of section 2.4.

2) The Sunyaev-Zeldovich effect is in principle a direct measure of the plasma column density. The angular resolution required for our analysis currently limits the use of this technique, however, future multi-frequency observations will determine independently both the temperature and number density profile of distant clusters purely through the S-Z effect (Hansen et al. 2002; Aghanim et al. 2003). The SZ-effect can thus be used to measure the electron density profile to large radii (SZ effect is proportional to n_e , whereas X-ray is proportional to n_e^2) and at large redshift (SZ is redshift independent).

3) Surface brightness: both from stellar light and radio observations of HI and molecular gas, one can in principle determine the baryon profile. An example is for M33 by (Corbelli 2003).

The distribution near black holes has been considered earlier. First by Peebles (1972) where energy consideration lead to the distribution $\rho \sim r^{-9/4}$, which was refined in (Bahcall & Wolf 1976) who found $\rho \sim r^{-7/4}$. Later numerical simulations have shown (Young 1980) that the profile near the black hole should be $\rho \sim r^{-3/2}$, which is just what we find. For accreting black holes this profile may be different (Freitag & Benz 2002).

4. Conclusion

We make a first attempt to derive analytically the asymptotic density profiles of baryonic structure in cosmology, which include galaxies, gaseous haloes and intra-clusters gas. We find that both disks and spherical solutions exist, and that generally the inner and outer density profile may be different. Thus we supply theoretical support for

the use of phenomenological profiles like

$$\rho_{\text{gas}}(r) = \frac{\rho(0)}{r^{\beta_1}(1+r)^{\beta_2}}. \quad (8)$$

For the disks we identify central cores with $\beta_d = 0$ from $\rho \sim r^\beta$. For the outer region we find $\beta_d = -3$. For spherical structures we identify both inner and outer profiles, which include $\beta_s = -6, -3, -3/2$. Our resulting profiles only apply to gaseous baryonic structures, but we point out a simple method whereby observations of the baryonic structure in principle allows one to deduce the dark matter density profile.

It is a pleasure to thank Tomas Bohr, James Binney, Sasha Dolgov, Lucio Mayer, Ben Moore and the people in Zurich for useful discussions, Victor Debattista and Marcella Carollo for observational guidance, and in particular Ruth Durrer for a valuable discussion.

REFERENCES

- Aghanim, N., Hansen, S. H., Pastor, S. & Semikoz, D. 2003, JCAP, 0305, 007
- Bahcall, J. N. & Wolf, R. A. 1976, ApJ, 209, 214
- Corbelli, E. 2003, preprint (astro-ph/0302318)
- Freitag, M. & Benz, W. 2002, Astronomy & Astrophysics, 394, 345
- Hansen, S. H., Horluck, S., Zauner, D., Dimon, P., Ellegaard, C. & Creagh, S. C. 1997, Phys. Rev. E, 55, 7048
- Hansen, S. H., Pastor, S. & Semikoz, D. V. 2002, ApJ, 573, L69
- Landau, L. D & Lifshitz E. M. 1987, Pergamon Press, UK
- Lewis, A. D., Buote, D. A. & Stocke, J. T. 2003, ApJ, 586, 135
- Moore, B., Quinn, T., Governato, F., Stadel, J. & Lake, G. 1999, MNRAS, 310, 1147
- Navarro, J. F., Frenk, C. S. & White, S. D. M. 1996, ApJ, 462, 563
- Peebles, P. J. E. 1972, ApJ, 178, 371
- Watanabe, S., Putkaradze, V. & Bohr, T. 2003, J. Fluid Mech., 480, 233
- Young, P. 1980, ApJ, 242,1232